

Optimization of Culture Condition in the Production of Bioenzymes by Bacteria Isolated from Poultry Waste in Sokoto State, Nigeria

¹Hauwa, H., ²S.B. Oyeleke and ³Adamu, Shahidah Ahmed

¹Faculty of Science, Department of Microbiology. Faculty of Science. Sokoto State University, Sokoto.

²Faculty of Science, Department of Microbiology. Federal University of Science And Technology Minna , Niger State.

³Department of Microbiology. Faculty of Science. Sokoto State University, Sokoto (08069303535) h_hauwau@yahoo.com

Abstract

Poultry wastes obtained from a poultry farm in Sokoto metropolis were analyzed for cellulose producing bacteria. *Bacillus megaterium*, *Bacillus laterosporus*, and *Bacillus amyloliquefeciens* isolated were screened for their ability to produce cellulase enzyme. All the isolates showed cellulase activity by exhibiting a wide halo on carboxymethylcellulase medium (CMC). The fermentation process was optimized using the following parameters, inoculum size, pH, Substrate concentration, temperature, and incubation periods. Cellulase activity was determined using DNS method, Banana peels was used as a substrate for the production of the enzymes, this was analysed with atomic absorption spectrophotometer (AAS). Cellulase enzyme was produced at inoculum size 1, 2, 3, 4, and 5%. pH 3, 5, 7, 9, and 11. Substrate concentration 1g, 2g, 3g, 4g, and 5g. Temperature 35, 45, 55, 65, and 75, for 1, 2, 3, 3, and 5, days respectively. *Bacillus laterosporus* recorded the highest cellulase activity of 0.37mg/ml in 5% substrate concentration among all the isolates while *Bacillus amyloliquefeciens* recorded highest cellulase production at pH3 with 45mg/ml *Bacillus laterosporus* recorded highest activity of cellulase production with 0.71mg/ml Temperature was also studied in the cellulase production and *Bacillus laterosporus* showed highest activity at 75°C with activity of 0.66mg/ml. This study showed that *Bacillus laterosporus* was the best cellulase producing bacteria among all the isolates.

Key words: poultry wastes, cellulase, Banana peels, Bacteria, Enzymes.

INTRODUCTION

Cellulose is the principal constituent of the cell wall of most terrestrial plants. The source of cellulose is in plants and it is found as microfibrils (2-20nm in diameter and 100 - 40,000nm long). These form the structurally strong frame work in the cell walls. Despite a worldwide and enormous utilization of natural cellulosic sources, there are still abundant quantities of cellulose-containing raw materials and waste products that are not exploited or which could be used more efficiently. Various biomass inducing residues including lignocellulosic material, paper waste, pulses, cereals straw and bagasses have been widely used as carbon sources for commercial cellulose fermentation Brijwani, and Vadlani, (2011), Wen, *et al.*, (2005) Belghith, *et al.*, (2001).

The problem in this respect is however to develop processes that are economically profitable. Cellulose-containing wastes may be agricultural, urban, or industrial in origin, sewage

sludge might also be considered a source of cellulose since its cellulosic content provides the carbon needed for methane production in the anaerobic digestion of sludge. Agricultural wastes include crop residue, animal excreta and crop processing wastes slashing generated in logging, saw dust formed in timber production and wood products in forestry originated activities (Pranner, 1979).

The previous negative attitude in which wastes were viewed self consciously as valueless and even offensive and for disposal only has been replaced in large part by a positive view in which wastes are recognized as raw materials of potential value (Pranner, 1979). Currently, there are two major ways of converting cellulose to glucose: chemical versus enzymatic. Enzymatic hydrolysis of cellulose is an important reaction in nature for it marks the first step in the decay of cellulose, the most abundantly occurring organic material.

In the early 1970s, the oil crisis generated interest in using cellulose as a chemical and energy resource. One promising approach was to hydrolyze the cellulose to glucose with fungal enzymes and then to ferment the glucose to ethanol which could be used as a liquid fuel (Mandels *et al.*, 1974).

Sources of Cellulase

Cellulases are easily obtained from microbial and fungal sources, but vertebrates lack the ability to produce endogenous cellulases, and hence are reliant upon gastrointestinal microorganisms for cell wall degradation of ingested plants. The beneficial effects of microorganisms in the digestive processes of terrestrial animals are well established (Combe *et al.*, 1976; McBee, 1977; Goldin, 1986; Moriarty, 1990). Some investigations have also suggested that microorganisms have a beneficial effect in fish digestive processes e.g. microbial breakdown of chitin (Minami *et al.*, 1972; Goodric and Morita, 1977; Danulat and Kausch, 1984; Kono *et al.*, 1987), p-nitrophenyl-B-N-acetylgalactosamine and collagen (MacDonald *et al.*, 1986), cellulose (Stickney and Shumway, 1974; Trust *et al.*, 1979; Saha and Ray, 1998; Bairagi *et al.*, 2002 a, b, 2004; Saha *et al.*, 2006), and the vitamin B12 producing ability of the bacteria (Sugita *et al.*, 1991). However, the specific cellulolytic activity shown by the bacterial species is found to depend on the source (Saxena *et al.*, 1993).

Microbial enzymes have the enormous advantage of being able to be produced in large quantities by established fermentation techniques. Enzyme production is closely controlled in micro organisms and therefore, to improve its productivity these controls can be exploited and modified. Cellulase yields appear to depend on a complex relationship involving a variety of factors like inoculum size (carbon source and cellulose quality), pH value, temperature, presence of inducers, medium additives, aeration, growth time, etc. (Immanuel *et al.*, 2006). Therefore, attention has been focused on studying the cellulolytic activity and cellulase enzyme production by several micro organisms in various products as well as in various environments. To establish a successful fermentation process it is necessary to make the environmental and nutritional conditions favorable for the micro organism for over-production of the desired metabolite.

Thus an elaborate investigation is therefore, required to establish the optimum conditions to scale up enzyme production in an individual fermentation process. Although there are reports

on the influence of various fermentation variables on cellulase production by different bacteria and fungi isolated from various natural environments (Garcia-Martinez *et al.*, 1980; Stewart and Parry, 1981; Coral *et al.*, 2002; Rajoka, 2004; Immanuel *et al.*, 2006), information on the optimum fermentation conditions for cellulase production by fish gut bacteria is lacking.

MATERIALS AND METHODS

The cellulolytic activity of the isolated organism was determined as described by Ghose, (1987). Loop full of each isolate was streaked aseptically on carboxymethyl cellulose medium. The plate was then incubated at 37 °C for 48 hours for bacterial isolates. After incubation period, the culture was flooded with 0.1% Congo red dye solution and later 40% NaCl solution and allowed to stand for 15-20 minutes. A zone of clearance formed around the bacterial colonies. The zone of clearance was measured using a meter rule. This represents the cellulolytic activity of the bacterial isolates.

Medium composition for cellulase production

The fermentation media used was Mary Mandels mineral salts solution and it was used along with different carbon and nitrogen sources. The medium (M1) contained the following (per L) Banana peels, 20g ; Peptone, 1g ; (NH₄)₂SO₄, 1.4g ; KH₂PO₄, 2g ; CaCl₂, 0.3g ; MgSO₄.7H₂O, 0.3g ; Urea, 0.3g ; trace metal solution (2.5g FeSO₄; 0.98g MnSO₄.H₂O ; 1.76g ZnSO₄.H₂O; 1.83g COCl₁₂.6H₂O dissolved in 1000ml of distilled water and heated to homogenized ; pH 6.8 (Jeffries, 1996).

Cellulase assay

Liquid media was used for the quantitative assay of cellulase production from the three bacterial strains. Cellulase activity was measured according to the method of Denison and Koehn (1977). The production of reducing sugar (glucose) from CMC substrate through cellulolytic activity was measured at 540 nm by the dinitrosalicylic acid method using glucose as the standard.

One cellulase unit (U) was defined as the amount of enzyme per milliliter culture filtrate that released 1 microgram glucose per minute.

Optimization of Cellulase Production Process

Effect of optimum pH on cellulase production by Bacteria isolates

The reaction mixtures of 120 hrs containing enzymes from the different bacteria isolates were prepared; the optimum pH for enzyme cellulase activity was examined by running the assay activity between pH ranges of 3.0, 5.0, 7.0, 9.0, and 11.

The enzyme activity for each pH was determined by adding DNSA reagents and the absorbance read at 540 nm.

Effect of optimum temperature on cellulase production by Bacteria isolates

The optimal temperature for activity was determined by assaying the activity of the enzyme at different temperature ranges of 35 °C, 45 °C, 55 °C, 65 °C, and 75 °C.

Effect of incubation period on cellulase production by Bacteria isolates

The incubation was carried out for different periods in this study including days 1, 2, 3, 4, and 5 for the bacteria isolates after which an assay was determined by Dinitrosalicylic method (Betrand *et al*, 2004).

Effect of inoculum size on cellulase production by Bacteria isolates

This activity was determined by assaying the enzymes at different sizes of 1%, 2%, 3%, 4% and 5% and the enzyme solution was maintained with different sizes of the inoculum.

Effect of substrate concentration on cellulase production by Bacteria isolates

The effect of substrate concentration was determined by assaying the activity of the enzymes of each bacteria with different substrate concentration of 1, 2, 3, 4 and 5%. DNSA reagent was added and the absorbance was taken on the spectrophotometer at 540nm as described by Betrand *et al.*, (2004).

RESULTS

The effect of substrate concentration on cellulase activity produced by bacterial isolates is shown in Fig1. *Bacillus laterosporus* had maximum activity recorded in 2% substrate concentration with cellulase activity of 0.35 mg/ml and the least activity recorded in 5% substrate concentration with activity of 0.15 mg/ml *Bacillus megaterium* recorded maximum cellulase activity in 2% substrate conc with activity of 0.37 mg/ml, which was then followed by a decrease in activity as the substrate concentration increases. *Bacillus amyloliquifeciens* recorded its least activity in 5% substrate concentration with activity of 0.14 mg/ml and the highest recorded at 1% substrate concentration with activity of 0.36 mg/ml.

The effect of pH on the activities of cellulase enzyme produced by bacterial isolates is shown in figure 2. From the result the highest activity produced by *Bacillus laterosporus* was at pH of 5 with activity of 0.36 mg/ml, these agrees with the report of Lynd *et al.*, 2002 and Gautam (2011) Who reported maximum yield at pH 5 and 6 by *A. niger* and *Trichoderma* sp while the least recorded at pH value of 9 with cellulase activity

of 0.10 mg/ml. *Bacillus megaterium* had an activity of 0.40 mg/ml at pH of 5 this was followed by a gradual decrease as the pH increases. *Bacillus amyloliquifeciens* recorded maximum activity of 0.45 mg/ml at pH of 3. There was a slight decrease as the pH approaches 11 with cellulase activity of 0.19 mg/ml. This shows that *Bacillus amyloliquifeciens* recorded the highest cellulase activity.

The effect of inoculum size on the activity of cellulase produced by bacterial isolates. It can be seen in figure 3 that *Bacillus laterosporus* exhibited high cellulase activity of 0.94 mg/ml and 0.65 mg/ml in 5% and 4% respectively, minimum cellulase activity was recorded at 3% with 0.46 mg/ml. *Bacillus megaterium* and *Bacillus amyloliquifeciens* also recorded maximum activity in 3%. *Bacillus megaterium* had cellulase activity of 0.85 mg/ml while *Bacillus amyloliquifeciens* had 0.89 mg/ml from the result *Bacillus laterosporus* recorded the highest cellulase activity while *Bacillus megaterium* recorded the least activity. Lower inoculum size require longer time for the cells to multiply to sufficient number to utilize the substrate and produce enzyme, an increase in the number of cells in the inoculums would ensure a rapid proliferation and biomass synthesis. When inoculum size was increased from 1-5% there was increases in enzyme production but after that the activity was decreased (Fig 3) due to depletion of nutrients by the enhanced biomass, which resulted dwindle in metabolic activity (Kashyap *et al.*, 2002). A balance between the increasing biomass and accessible nutrient would yield an optimal enzyme production (Ramachandran *et al.*, 2004).

The effect of incubation period on the activity of cellulase producing isolates. From figure 4 it can be seen that *Bacillus laterosporus* exhibited high cellulase activity of 0.71 mg/ml after 120 hrs (day 5) These agrees with the report of Gautam *et al.*, (2011) who reported optimum yield on the 5th day but contrary to his other report of highest cellulase activity by *A.niger* and *Trichoderma* sp on the 4th and 6th day which was suitable for commercial point of view Kang *et al.*, (2004) , and it might be due to the depletion of nutrients in the medium which stressed the fungal physiology resulting in the inactivation on secretory machinery of the enzymes (Nochur *et al.*, 1993), the least activity recorded was at 48 hrs (day 2) with 0.4 mg/ml. *Bacillus megaterium* recorded its highest activity after 48 hrs (day 2) (0.56 mg/ml) this was followed by a decline in cellulase activity at 72 hrs (day 3) and 120 hrs (day 5).

The least activity recorded was at 96 hrs (day 4) (0.21 mg/ml) *Bacillus amyloliquifeciens* had its highest activity recorded at 24 hrs (day 1) this was followed by a slight decrease as the hours increases, to 120 hrs (day 5) the activity drop to 0.18 mg/ml which was the least recorded.

The effect of temperature on cellulase production by the bacterial isolates is shown in figure 5 from the figure *Bacillus laterosporus* had cellulase activity of 0.04 mg/ml at 35 °C, this was followed by a slight increase at 45 °C (0.55 mg/ml) these disagrees with the report of Ray *et al* (2007) who reported that minimum cellulose yield was observed when fermentation was carried out at 45 °C by *B. subtilis* and *B.circulans* it was then followed by a sharp decrease at 55°C and 65°C 0.23 mg/ml and 0.13 mg/ml but as the temperature increases to 75°C the activity also increases to 0.66 mg/ml. *Bacillus megaterium*

exhibited its high cellular activity at 55°C with 0.58 mg/ml but the activities fluctuated by increasing and decreasing as temperature rises and fall the lowest recorded for *Bacillus megaterium* was at 35°C with cellulase activity of 0.36 mg/ml . *Bacillus amyloliquifeciens* had its highest activity at 35°C with 0.59 mg/ml, it was then followed by a slight decrease as the temperture increases with the lowest activity at 75°C (0.30 mg/ml). From the result *Bacillus laterosporus* recorded the highest cellulase activity. These agrees with the findings of Bakare *et al.*, (2005) who found that cellulose enzyme produced by *pseudomonas fluorescence* was activated at 30-35 °C, also disagrees with the report of immanuel *et al.*, 2006 who reported maximum endogluconase activity in *cellulomonas Bacillus* and *Micrococcus* sp at 40 °C .




Figure 1; Effect of substrate concentration on cellulase activity produced by *B. laterosporus*, *B. megaterium*, *B. amyloliquifeciens*




Figure2: Effect of pH on cellulase activity produced by *B. laterosporus*, *B. megaterium*, *B. amyloliquifeciens*




Figure 3: Effect of inoculum size on cellulase activity produced by *B. laterosporus*, *B. megaterium*, *B. amyloliquifeciens*




Figure 4: Effect of incubation periods on cellulase activity produced by *B. laterosporus*, *B. megaterium*, *B. amyloliquifeciens*




Figure 5: Effect of Temperature on cellulase activity produced by *B. laterosporus*, *B. megaterium*, *B. amyloliquifeciens*

DISCUSSION

Three species of *Bacillus* were characterized and identified on the basis of macroscopy and microscopy. The macroscopic and microscopic characterization revealed the presence of *Bacillus laterosporus*, *Bacillus megaterium*, and *Bacillus amyloliquifeciens*. All the bacterial isolates were hydrolizers of carboxymethyl cellulose medium, based on cellulolytic activity of the isolates following their zones of clearing on CMC medium. Temperature and pH are the most important factor that influence enzyme activity. *Bacillus laterosporus* had an enzyme activity of 0.44 mg/ml at 35°C this was followed by an increase in enzyme activity as the temperature increase, with maximum activity of 0.56 mg/ml. cellulase enzymes when hydrolyzed by cellulolytic microorganisms instead of being left alone for natural degradation, these enzymes can be utilized effectively under optimum conditions, to produce cellulase as realized as the temperature increased beyond 45°C the enzyme activity decreased, The minimum enzyme activity for *Bacillus laterosporus* was recorded at 75°C with enzyme activity of 0.17 mg/ml while *Bacillus magaterium* had an enzyme activity of 0.56 mg/ml at 35°C and the temperature increased to 45°C enzyme activity also increases to 0.66 mg/ml this was followed by a sharp increase in enzyme activity of 0.82

mg/ml at 55°C. Hence, further increase in temperature lead to a decrease in enzyme activity of 0.32 mg/ml at 75°C. This is in line with the report of Kelly *et al* (1997) that during isomerization, temperature is preferably maintained within the range of 20-90°C and the best activity is obtained at 50-75°C.

Further increase in cellulose concentration beyond the level that gave optimum cellulase enzyme production will result in proportionate decrease in yield. This is in variance with Haapela *et al.*, (1995) and Jeffries, (1996) who reported that maximum enzyme activity was recorded with the medium concentration of cellulose at 10g/l. Lynd *et al.*, (2002), reported that the production of cellulase for the utilization of cellulose is induced only in the presence of specific substrate but suppressed when easily utilizable sugars such as glucose are available.

CONCLUSION

This study revealed that poultry wastes and Banana peel, are of domestic and industrial agrowastes, that can be use to produce large amounts of cellulase enzymes when hydrolyzed by cellulolytic microorganisms instead of being left alone for natural degradation, these enzymes can be utilized effectively under optimum conditions, to produce cellulase.

REFERENCES

- Bairagi A., Sarkar Ghosh K., Sen S.K., and Ray A.K. (2002a). Enzyme producing bacterial flora isolated from fish digestive tracts. *Aquaculture International* **10**: 109-121.
- Bakare, M. K., Adewale I.O. Ajayi A. and Shonukan O.O. (2005). Purification and Characterization of Cellulase from the wild-type and two Improved Mutants of *Pseudomonas fluorescens*, *African journal of biotechnology*, Vol.4 (9) pp 898-904
- Belghith, H., Ellouz-Chaabouni, S. and Gargouri, A. (2001). Stabilization of *Penicillium occitanis* Cellulases by Spray Drying in Presence of Maltodextrin. *Enzyme and Microbial Technology*, **28**, 253-258.
- Bertrand, T.F., Fredrie, T. and Robert, N. (2004). Production and partial characterisation of a thermostable amylase from *Ascomycetes yeast* strain isolated from starchy soil. Mc Graw Hill inc. New York. PP 57-59.
- Brijwani, K. and Vadlani, P.V. (2011). Cellulolytic Enzymes Production via Solid State Fermentation: Effect of Pretreatment Methods on Physiochemical Characteristics of Substrate. *Enzyme Research*, Vol 86 :134.
- Combe E., Dermanne Y., Gueguen L., Ivorec-Szylit O., Meslin J.H., and Sacquet E. (1976). Some aspects of the relationships between gastrointestinal flora and host nutrition. *World Review of Nutrition and Dietetics* **24**: 1-57.
- Coral, G. Arian, B., Unaldi M.N. and Guvenmez, H. (2002). Thermostable alkaline protease produced by an *Aspergillus niger* strain. *Journals of Microbiology*. **53**:491-498.
- Danulat E., and Kausch, H. (1984). Chitinase activity in the digestive tract of the *Cod*, *Gadus morhua* (L.). *Journal of Fish Biology* **24**: 125-133.
- Denison, D.A. and Koehn, R.D. (1977). Cellulase activity of *pronia Oedipus*. *Mycologia*, **69**:592-603.
- Garcia-Martinez, D.V. Shimuyo A., Madia A., and Deman A.L., (1980). Studies on cellulase production by *Clostridium Thermocellum*. *European Journal Applied Microbiology and Biotechnology* **9**: 189-197.

- Gauam, S.P. Bundela, P, S. Pandey, A.K. Jamaluddin Khan, Awasthin M.K. and Sarsanya S. (2011). Optimization for the production of cellulose enzyme from municipal solid waste residue by two novel cellulolytic fungi. *Biotechnology research international*. Vol. 2:11 Article ID 810428.PP8
- Goldin B.R. (1986). *In situ* bacterial metabolism and colon mutagens. *Annual Review of Microbiology* **40**: 367-393.
- Goodrich.D., and Morita R.Y. (1977). Bacterial chitinase in the stomach of marine fishes from Yaquina Bay, Oregon, USA. *Marine Biology* **41**: 355-360.
- Haapela, R., Parkkinen, E., Susminen, P., Unk [1] Abe J, Bergman FW, Obata K, and Hikuri S (1988). Production of raw starch digesting amylase by *Aspergillus* K-27. *Applied. Microbiology. Biotechnoogy*. **27**(1): 447- 450.
- Immanuel G., Dhanusha R., Prena P., and Palavesan A. (2006) Effect of different growth parameters on endoglucanase enzyme activity by bacteria isolated from coir retting effluents of estuarine environments. *International Journal of environmental Science and Technology*. **3**: 25-34.
- Jeffries, T.W. (1996). Production and Applications of Cellulase Laboratory Procedures, 1-10.
- Kang S.W., Park Y.S., Lee J.S., Hong SI and Kim S.W. Production of cellulases and Hemicellulases by *Aspergillus niger* KK2 from lignocellulosic biomass. *BioresourceTechnology* Vol. 91920 pp153-156.
- Kasyap, P., Sabu A., Pandey A, and Szakacs G. (2002) Extracellular leglutaminase production by *Zygosaccharomyces rouxii* under solid state fermentation. *process Biochem* **38**; 307-312.
- Kelly, C.T., M.A., Tigue., E.M., Doyle and Fogarty M.M.(1997). Raw Starch degrading alkaline amylase of *Bacillus* SP. *Journal of Industrial Microbiology.*, **15** : 446-448.
- Kono, M., Matsui, T., and Shimizu C. (1987). Chitin-decomposing bacteria in digestive tracts of cultured red sea bream and Japanese eel. *Bulletin of the Japanese Society of Scientific Fisheries* (Nippon Suisan Gakkaishi) **53**: 305-310.
- Lynd, L.R., Weimer P.J., Vanzyl W.H. and Pretorius I.S. (2002). Microbial cellulose utilization: mixed culture of *Aspergillus niger* and *Saccharomyces cerevisiae* grown on Sorghum. *Microbial molecular. Biogyl Review*. **66**(3): 506-577.
- Mandels, M., Hontz, L. and Nyostrom, J. (1974). Enzymatic hydrolysis of Waste Cellulose *Biotechnology Bioeng* **16**: 1471-1493.
- McBee R.H., Clarke R.T.J., and Bauchop T. (1977). Fermentation in the hindgut. (eds.) *Microbial ecology of the gut*. Academic Press, London. Pp.185-222
- Minami, Z., Kajinta, M., and Hibino S. (1972). Studies on the utilization of petro-yeast as diet of fresh water fish. The distribution of chitin-decomposing bacteria in digestive tracts. *Bulletin of Japanese Society of Scientific Fisheries* **38**: 1013-1019.
- Moriarty, D.J.W. (1990) Interactions of microorganisms and aquatic animals, particularly the nutritional role of the gut flora., In; Lesel R. (ed) *Microbiology in poikilotherms*. Elsevier, Amsterdam. pp 217-222.
- Nochur S.V., Robertsm, F. and Deman A.I. True cellulose production by clostridium grown on different carbon source. *Journal of bacteriology* Vol 15 6 pp 641-646.
- Pranner, J. (1979). Environmental Microbiology and Waste Utilization G//AMV Proceedings (ed emejuaiwe), Academic Press, London, 67-69.
- Rajoka. (2004). Influence of various fermentation variables on exo-glucanase production in *cellulomonas flavigena*. *Electronic Journal of Biotechnology* **43**:15 22.
- Ramachandran, S., Patel A.K., Nanpiothiri, K. M., Francis F., Nagy V., Szakac S.G., Pandey A. (2004). Coconut oil cake a potential raw material for the production of amylase. *Bioresource Technol* **93**;169–174.
- Ray, A.K., Bairagi A, Sarkar Ghosh k, and Sen S.K. (2007). Optimization of fermentation conditions for cellulose production by *Bacillus subtilis* CY5 and *Bacillus circulans* TP3. Isolated from Fish Gut, *Acta ichthyologicaet piscatorial*, Vol **37** (1) pp 47-53.
- Saha A.K., and Ray A.K. (1998). Cellulase activity in rohu fingerlings. *Aquaculture International* **6**: 281-291.
- Saxena S., Bahadur J.,and Varma A. (1993). Cellulose and hemi-cellulose degrading bacteria from termite gut and mound soils of India. *International Journal of Microbiology* **33**: 55-60.

- Stewart J.C, and Parry J.B. (1981). Factors influencing the production of cellulase by *Aspergillus fumigates* (Fresenius). *Journal of General Microbiology* **125**: 33-39.
- Stickney R R., and Shunway S.E. (1974). Occurance of cellulose activity in the stomachs of fish, *Journal of fish Biology* **6**:779-790.
- Sugita H., Miyajima C., and Deguchi Y. (1991). The vitamin B12-producing ability of the intestinal microflora of freshwater fish. *Aquaculture* **92**: 267-276.
- Trust T.J., Bull L.M., Currie B.R., and Buckley J.T. (1979). Obligate anaerobic bacteria in the gastrointestinal microflora of the grass carp. (*Ctenopharyngodon idella*), goldfish. (*Carassius auratus*), and rainbow trout (*Salmo gairdneri*). *Journal of Fisheries Research Board of Canada* **36**:1174-1179.
- Wen, Z., Liao, W. and Chen, S. (2005). Production of Cellulase/ β -Glucosidase by the Mixed Fungi Culture *Trichoderma reesei* and *Aspergillus phoenicis* on Dairy Manure. *Process Biochemistry*, **40**, 3087-3094.